

論文発表

- 1) 生理活性脂質「環状ホスファチジン酸」の生理作用とその活用
嶋田有紀子 室伏きみ子 Journal of Beauty & Wellness Research 第1巻 10-18 (2025)
- 2) 美容系職業を目指す学生における自己と職業に対するジェンダー意識に関する予備調査
横田 悠季 猪崎 弥生 嶋田 有紀子 Journal of Beauty & Wellness Research 第1巻 78-83 (2025)
- 3) A novel sphingomyelin/cholesterol domain-specific probe reveals the dynamics of the membrane domains during virus release and in Niemann-Pick type C.
Makino A., Abe M., Ishitsuka R., Murate M., Kishimoto T, Sakai S., Hullin-Matsuda F., Shimada Y., Inaba T., Miyatake H., Tanaka H., Kurahashi A., Pack C.G., Kasai R.S., Kubo S., Schieber N.L., Dohmae N., Tochio N., Hagiwara K., Sasaki Y., Aida Y., Fujimori F., Kigawa T., Nishibori K., Parton R.G., Kusumi A., Sako Y., Anderluh G., Yamashita M., Kobayashi T., Greimel P, Kobayashi T
FASEB J. ;31(4):1301-1322 (2017)..
- 4) Translation of lysosomal cathepsin D caused by oxidative stress or proteasome inhibition in primary cultured neurons and astrocytes.
Miura Y., Sakurai Y., Hayakawa M., Shimada Y., Zempel H., Sato Y., Hisanaga S., Endo T.
Biol. Pharm. Bull., 33, 22-28, (2010)
- 5) Plasma membrane microdomains in aging and disease.
Ohno-Iwashita Y, Shimada Y, Hayashi M, Inomata M
Geriatr Gerontol Int. 10 Suppl 1:S41-52 (2010).
- 6) Cholesterol-binding toxins and anti-cholesterol antibodies as structural probes for cholesterol localization.
Ohno-Iwashita Y, Shimada Y, Hayashi M, Iwamoto M, Iwashita S, Inomata M.
Subcell Biochem. 51:597-621(2010).
- 7) Age-associated up-regulation of a negative co-stimulatory receptor PD-1 in mouse CD4+ T cells.
Shimada Y., Hayashi M., Ohno-Iwashita Y., Inomata M.
Exp. Gerontology, 44, 417-522, (2009)
- 8) MOV10 as a novel telomerase-associated protein.
Nakano M, Kakiuchi Y, Shimada Y, Ohyama M, Ogiwara Y, Sasaki-Higashiyama N, Yano N, Ikeda F, Yamada E, Iwamatsu A, Kobayashi K, Nishiyama K, Ichikawa S, Kaji K, Ide T, Murofushi H, Murakami-Murofushi K.
Biochem Biophys Res Commun. 388, 328-32, (2009)
- 9) Development of SHIP2 transgenic mice and determination of their phenotype.
Shimada Y, Ohta M, Kanai S, Kawakami T, Takiguchi A, Funakoshi A, Sekime A, Mita K, Miyasaka, K
Digestion & Absorption, 31, 91-96, (2008)
- 10) Impairment in a negative regulatory system for TCR signaling in CD4+ T cells from old mice.
Inomata M, Shimada Y, Hayashi M, Shimizu J, Ohno-Iwashita Y.
FEBS Lett. 581, 3039-3043, (2007)

- 11) Increased lipid rafts and accelerated lipopolysaccharide-induced tumor necrosis factor- α secretion in Abca1-deficient macrophages.
Koseki M, Hirano K, Masuda D, Ikegami C, Tanaka M, Ota A, Sandoval JC, Nakagawa-Toyama Y, Sato SB, Kobayashi T, Shimada Y, Ohno-Iwashita Y, Matsuura F, Shimomura I, Yamashita S.
J Lipid Res. 48, 299-306, (2007)
- 12) Detection of cholesterol-rich microdomains in the inner leaflet of the plasma membrane.
Hayashi M, Shimada Y, Inomata M, Ohno-Iwashita Y.
Biochem Biophys Res Commun. 351, 713-718, (2006)
- 13) Detachment-associated changes in lipid rafts of senescent human fibroblasts.
Inomata M, Shimada Y, Hayashi M, Kondo H, Ohno-Iwashita Y.
Biochem Biophys Res Commun. 343, 489-495, (2006)
- 14) Separation of a cholesterol-enriched microdomain involved in T-cell signal transduction.
Shimada Y, Inomata M, Suzuki H, Hayashi M, Abdul Waheed A, Ohno-Iwashita Y.
FEBS J. 272, 5454-5463, (2005)
- 15) Structure and function of lipid rafts in human activated T cells.
Tani-ichi S, Maruyama K, Kondo N, Nagafuku M, Kabayama K, Inokuchi J, Shimada Y, Ohno-Iwashita Y, Yagita H, Kawano S, Kosugi A.
Int Immunol. 17, 749-758, (2005)
- 16) Distinct localization of lipid rafts and externalized phosphatidylserine at the surface of apoptotic cells.
Ishii H, Mori T, Shiratsuchi A, Nakai Y, Shimada Y, Ohno-Iwashita Y, Nakanishi Y.
Biochem Biophys Res Commun. 327, 94-99, (2005)
- 17) Perfringolysin O, a cholesterol-binding cytolysin, as a probe for lipid rafts.
Ohno-Iwashita Y, Shimada Y, Waheed AA, Hayashi M, Inomata M, Nakamura M, Maruya M, Iwashita S.
Anaerobe. 10, 125-134, (2004)
- 18) Axon-dominant localization of cell-surface cholesterol in cultured hippocampal neurons and its disappearance in Niemann-Pick type C model cells.
Tashiro Y, Yamazaki T, Shimada Y, Ohno-Iwashita Y, Okamoto K.
Eur J Neurosci. 20, 2015-2021., (2004)
- 19) Accelerated Abeta aggregation in the presence of GM1-ganglioside -accumulated synaptosomes of aged apoE4-knock-in mouse brain.
Yamamoto N, Igbabvoa U, Shimada Y, Ohno-Iwashita Y, Kobayashi M, Wood WG, Fujita SC, Yanagisawa K.
FEBS Lett. 569, 135-139, (2004)
- 20) Reduced sensitivity of Niemann-Pick C1-deficient cells to theta-toxin (perfringolysin O): sequestration of toxin to raft-enriched membrane vesicles.
Ohsaki Y, Sugimoto Y, Suzuki M, Kaidoh T, Shimada Y, Ohno-Iwashita Y, Davies JP, Ioannou YA, Ohno K, Ninomiya H.
Histochem Cell Biol. 121, 263-272, (2004)

- 21) A novel cholesterol stain reveals early neuronal cholesterol accumulation in the Niemann-Pick type C1 mouse brain. Reid PC, Sakashita N, Sugii S, Ohno-Iwashita Y, Shimada Y, Hickey WF, Chang TY. *J Lipid Res.* 45, 582-591, (2004)
- 22) Gamma-secretase activity is present in rafts but is not cholesterol -dependent.
Wada S, Morishima-Kawashima M, Qi Y, Misono H, Shimada Y, Ohno-Iwashita Y, Ihara Y. *Biochemistry.* 42, 13977-13986, (2003)
- 23) Cellular aging-dependent decrease in cholesterol in membrane microdomains of human diploid fibroblasts.
Nakamura M, Kondo H, Shimada Y, Waheed AA, Ohno-Iwashita Y. *Exp Cell Res.* 290, 381-390, (2003)
- 24) Reduction of glycosphingolipid levels in lipid rafts affects the expression state and function of glycosylphosphatidylinositol-anchored proteins but does not impair signal transduction via the T cell receptor.
Nagafuku M, Kabayama K, Oka D, Kato A, Tani-ichi S, Shimada Y, Ohno-Iwashita Y, Yamasaki S, Saito T, Iwabuchi K, Hamaoka T, Inokuchi J, Kosugi A. *J Biol Chem.* 278, 51920-51927, (2003)
- 25) Recycling compartments and the internal vesicles of multivesicular bodies harbor most of the cholesterol found in the endocytic pathway.
Möbius W, van Donselaar E, Ohno-Iwashita Y, Shimada Y, Heijnen HF, Slot JW, Geuze HJ. *Traffic.* 4, 222-231, (2003)
- 26) Ultrastructural localization of flotillin-1 to cholesterol-rich membrane microdomains, rafts, in rat brain tissue.
Kokubo H, Helms JB, Ohno-Iwashita Y, Shimada Y, Horikoshi Y, Yamaguchi H. *Brain Res.* 965, 83-90, (2003)
- 27) Biotinylated theta-toxin derivative as a probe to examine intracellular cholesterol-rich domains in normal and Niemann-Pick type C1 cells.
Sugii S, Reid PC, Ohgami N, Shimada Y, Maue RA, Ninomiya H, Ohno-Iwashita Y, Chang TY. *J Lipid Res.* 44, 1033-1041, (2003)
- 28) The C-terminal domain of perfringolysin O is an essential cholesterol -binding unit targeting to cholesterol-rich microdomains.
Shimada Y, Maruya M, Iwashita S, Ohno-Iwashita Y. *Eur J Biochem.* 269, 6195-6203, (2002)
- 29) Lipid binding activity of a neuron-specific protein NAP-22 studied in vivo and in vitro.
Terashita A, Funatsu N, Umeda M, Shimada Y, Ohno-Iwashita Y, Epand RM, Maekawa S. *J Neurosci Res.* 70, 172-179, (2002)
- 30) Immunoelectron microscopic localization of cholesterol using biotinylated and non-cytolytic perfringolysin O.
Möbius W, Ohno-Iwashita Y, van Donselaar EG, Oorschot VM, Shimada Y, Fujimoto T, Heijnen HF, Geuze HJ, Slot JW. *J Histochem Cytochem.* 50, 43-55, (2002)
- 31) Selective binding of perfringolysin O derivative to cholesterol-rich membrane microdomains (rafts).

- Waheed AA, Shimada Y, Heijnen HF, Nakamura M, Inomata M, Hayashi M, Iwashita S, Slot JW, Ohno-Iwashita Y.
Proc Natl Acad Sci U S A. 98, 4926-4931, (2001)
- 32) Contribution of histidine residues to oligomerization of theta-toxin (perfringolysin O), a cholesterol-binding cytolysin.
Nakamura M, Sekino-Suzuki N, Shimada Y, Ohno-Iwashita Y.
Biosci Biotechnol Biochem. 63, 1640-1643, (1999)
- 33) The conserved undecapeptide shared by thiol-activated cytolysins is involved in membrane binding.
Jacobs T, Cima-Cabal MD, Darji A, Méndez FJ, Vázquez F, Jacobs AA, Shimada Y, Ohno-Iwashita Y, Weiss S, de los Toyos JR.
FEBS Lett. 459, 463-466, (1999)
- 34) Cross-linking of plasmalemmal cholesterol in lymphocytes induces capping, membrane shedding, and endocytosis through coated pits.
Hagiwara H, Kogure SY, Nakamura M, Shimada Y, Ohno-Iwashita Y, Fujimoto T.
Biochem Biophys Res Commun. 260, 516-521, (1999)
- 35) C-terminal amino acid residues are required for the folding and cholesterol binding property of perfringolysin O, a pore-forming cytolysin.
Shimada Y, Nakamura M, Naito Y, Nomura K, Ohno-Iwashita Y.
J Biol Chem. 274, 18536-18542, (1999)
- 36) A novel 66-kDa stress protein, p66, associated with the process of cyst formation of *Physarum polycephalum* is a *Physarum* homologue of a yeast actin-interacting protein, AIP1.
Matsumoto S, Ogawa M, Kasakura T, Shimada Y, Mitsui M, Maruya M, Isohata M, Yahara I, Murakami-Murofushi K.
J Biochem. 124, 326-331, (1998)
- 37) Cell cycle-dependent activation of telomerase in naturally synchronized culture of a true slime mold, *Physarum polycephalum*.
Shimada Y, Nakano M, Kanda N, Murakami-Murofushi K, Kim JK, Ide T, Murofushi H.
Biochem Biophys Res Commun. 232, 492-496, (1997)
- 38) Telomere and telomerase in human leukocytes and immortal cell lines.
Shimada Y, Nakano M, Murakami-Murofushi K
Nat. Sci. Ochanomizu Univ. 47, 35-43, (1996)
- 39) Reduced fidelity of DNA synthesis in cell extracts from chemically induced primary thymic lymphomas of mice.
Kubota Y, Murakami-Murofushi K, Shimada Y, Ogiu T, Oikawa T.
Cancer Res. 55, 3777-3780, (1995)
- 40) Isolation of a new species of *Physarum* lysophosphatidic acid, PHYLPA, and its effect on DNA polymerase activity.
Takahashi Y, Shimada Y, Shioda M, Yoshida S, Murofushi H, Murakami-Murofushi K.
Cell Struct Funct. 18, 135-138, (1993)
- 41) Expression of a 66-kD heat shock protein associated with the process of cyst formation of a true

slime mold, *Physarum polycephalum*.

Shimada Y, Kasakura T, Yokota M, Miyata Y, Murofushi H, Sakai H, Yahara I, Murakami-Murofushi K.

Cell Struct Funct. 17, 301-309, (1992)

- 42) Changes in preference for receptor subtypes of configurational variants of a glutamate analog: conversion from the NMDA-type to the non-NMDA type.

Ishida M, Ohfuné Y, Shimada Y, Shimamoto K, Shinozaki

Brain Res. 550, 152-156, (1991)

- 43) Extracellular polysaccharide from a true slime mold, *Physarum polycephalum*. Polymerization and autodegradation associated with cell growth, and its inhibitory activity on eukaryotic DNA polymerase.

Shimada Y, Murakami-Murofushi K, Shioda M Ohta J

Nat. Sci. Ochanomizu Univ. 42, 39-47, (1991)

- 44) Changes in phospholipid composition and phospholipase D activity during the differentiation of *Physarum polycephalum*.

Minowa A, Kobayashi T, Shimada Y, Maeda H, Murakami-Murofushi K, Ohta J, Inoue K.

Biochim Biophys Acta. 1043, 129-133, (1990)